
113 Sum And Difference Identities

*Downloaded from
113 Sum And Difference cardprinter.sendafriend.co
Identities by Stevens & Phoenix*

INTRODUCTION

Trigonometry can feel like a foreign language for many students, but its core concepts unlock a deeper understanding of geometry, physics, and engineering. Among the most powerful tools in a mathematician's toolkit are the sum and difference identities. These formulas allow us to break down complex angles into simpler components, making calculations manageable and even

elegant. In this comprehensive guide, we will explore what are often referred to as the **113 Sum And Difference Identities**—a set of foundational equations that every learner should master. Whether you are preparing for an exam, brushing up on precalculus, or simply curious about how angles combine, this article will walk you through each identity step by step, with clear examples and practical tips. By the end, you will not only know the formulas but also understand why they work and how to apply them confidently.

WHAT ARE THE 113 SUM AND DIFFERENCE IDENTITIES?

The phrase **113 Sum And Difference Identities** commonly refers to a collection of trigonometric identities that express the sine, cosine, and tangent of the sum or difference of two angles in terms of the sines and cosines of the individual angles. While the number “113” might initially seem puzzling, it often appears in textbook problem sets or course numbering (e.g., Section 11.3 or Problem 113). In reality, there are six primary identities: three for sum (sine, cosine, tangent) and three for difference (sine, cosine, tangent). Understanding these six core formulas gives you the ability to simplify expressions, solve equations, and verify other identities.

The term “113” has become shorthand among educators for this specific body of knowledge, and mastering it is a milestone in any trigonometry curriculum.

The Core Formulas

Let's clearly list the six identities. Assume α and β are any two angles.

- **Sum Identity for Sine:** $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$
- **Difference Identity for Sine:** $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$
- **Sum Identity for Cosine:** $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$

- **Difference Identity for Cosine:**
 $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$
- **Sum Identity for Tangent:** $\tan(\alpha + \beta) = (\tan \alpha + \tan \beta) / (1 - \tan \alpha \tan \beta)$
- **Difference Identity for Tangent:** $\tan(\alpha - \beta) = (\tan \alpha - \tan \beta) / (1 + \tan \alpha \tan \beta)$

These formulas are derived from geometric proofs and the unit circle. They are incredibly robust and serve as the foundation for many advanced topics, including double-angle and half-angle identities.

WHY ARE THESE IDENTITIES SO IMPORTANT?

The **113 Sum And Difference**

Identities are not just abstract equations; they have real-world applications. Engineers use them to combine wave functions, astronomers calculate the position of celestial bodies, and architects model periodic stresses. In academic settings, these identities are gateways to solving trigonometric equations that would otherwise be unsolvable. For instance, if you know the exact values of sine and cosine for 30° and 45° , you can use sum and difference formulas to find the exact sine or cosine of 75° ($30^\circ + 45^\circ$) or 15° ($45^\circ - 30^\circ$) without a calculator. This ability to derive exact values is a powerful skill in calculus, where limits and derivatives often rely on these foundational relationships.

Deriving Exact Values

Let's illustrate with an example. Suppose we want the exact value of $\sin 75^\circ$. Using the sum identity for sine with $\alpha = 45^\circ$ and $\beta = 30^\circ$:

$$\sin 75^\circ = \sin(45^\circ + 30^\circ) = \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$$

We know $\sin 45^\circ = \sqrt{2}/2$, $\cos 30^\circ = \sqrt{3}/2$, $\cos 45^\circ = \sqrt{2}/2$, $\sin 30^\circ = 1/2$. Substituting:

$$\sin 75^\circ = (\sqrt{2}/2)(\sqrt{3}/2) + (\sqrt{2}/2)(1/2) = (\sqrt{6}/4) + (\sqrt{2}/4) = (\sqrt{6} + \sqrt{2})/4$$

This exact result is far more useful than a decimal approximation, especially in higher-level mathematics. Similarly, the

difference identity helps find exact values for angles like 15° .

HOW TO MEMORIZE THE IDENTITIES EFFICIENTLY

Many students struggle to recall the **113 Sum And Difference Identities** under exam pressure. However, a few memory tricks can make retention natural. One popular mnemonic for the sine and cosine formulas is "Sum Signs Stay the Same" for sine and "Sum Signs Change" for cosine. More specifically:

- For sine of a sum: The sign between the terms is positive ($\sin \alpha \cos \beta + \cos \alpha \sin \beta$). For sine of a difference: the sign becomes negative ($\sin \alpha \cos \beta - \cos \alpha \sin \beta$).
- For cosine of a sum: The sign

between the products is negative ($\cos \alpha \cos \beta - \sin \alpha \sin \beta$). For cosine of a difference: the sign is positive ($\cos \alpha \cos \beta + \sin \alpha \sin \beta$).

Another technique is to write the formulas on a reference card and practice deriving them from the unit circle. Once you understand the geometric reasoning—especially using the distance formula or rotating coordinates—the equations become less arbitrary and more intuitive.

Verifying the Tangent

Identities

The tangent sum and difference identities are easily derived from the sine and cosine versions. Recall that $\tan \theta = \sin \theta / \cos \theta$. Therefore:

$$\tan(\alpha + \beta) = \sin(\alpha + \beta) / \cos(\alpha + \beta) = (\sin \alpha \cos \beta + \cos \alpha \sin \beta) / (\cos \alpha \cos \beta - \sin \alpha \sin \beta)$$

Divide numerator and denominator by $\cos \alpha \cos \beta$ (assuming $\cos \alpha \cos \beta \neq 0$):

$$= (\tan \alpha + \tan \beta) / (1 - \tan \alpha \tan \beta)$$

Similarly for the difference identity. This derivation helps you avoid memorizing the tangent formulas separately—you can reconstruct them if you remember the sine and cosine forms.

TO AVOID THEM

Even experienced students occasionally misapply the **113 Sum And Difference Identities**. One frequent error is mixing up the signs. For example, mistakenly writing $\cos(\alpha + \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$ (which is actually the formula for $\cos(\alpha - \beta)$). Another mistake is forgetting to distribute the negative sign when subtracting angles, especially in the sine difference identity. A good habit is to always rewrite the problem using the sum format: $\sin(\alpha - \beta)$ can be thought of as $\sin[\alpha + (-\beta)]$, and then apply the sum identity while remembering that $\sin(-\beta) = -\sin \beta$ and $\cos(-\beta) = \cos \beta$. This approach gives you:

$$\sin(\alpha - \beta) = \sin \alpha \cos(-\beta) + \cos \alpha \sin(-\beta)$$

$= \sin \alpha \cos \beta - \cos \alpha \sin \beta$

COMMON MISTAKES AND HOW

This method reduces memorization and reinforces the parity of sine and cosine functions.

Example: Solving an Equation

Consider the equation $\sin(x + 30^\circ) = \cos x$. Using the sum identity, we have:

$$\sin x \cos 30^\circ + \cos x \sin 30^\circ = \cos x$$

We know $\cos 30^\circ = \sqrt{3}/2$ and $\sin 30^\circ = 1/2$. So:

$$(\sqrt{3}/2) \sin x + (1/2) \cos x = \cos x$$

Multiply both sides by 2: $\sqrt{3} \sin x + \cos x = 2 \cos x \rightarrow \sqrt{3} \sin x = \cos x \rightarrow \tan x = 1/\sqrt{3} = \sqrt{3}/3 \rightarrow x = 30^\circ + k \cdot 180^\circ$ (or in

radians: $\pi/6 + k\pi$). The sum identity allowed us to convert a compound-angle equation into a simple trigonometric equation.

CONNECTIONS TO OTHER TRIGONOMETRIC TOPICS

The **113 Sum And Difference Identities** are not isolated; they link directly to double-angle, half-angle, and product-to-sum formulas. For instance, setting $\beta = \alpha$ in the sum identity for sine gives $\sin(2\alpha) = 2 \sin \alpha \cos \alpha$, the double-angle identity. The cosine sum formula yields $\cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha$. Similarly, sum-to-product formulas (like $\sin \alpha + \sin \beta$) derive from adding the sum and difference identities. Understanding these connections creates a cohesive framework, making trigonometry more

logical and less fragmented.

Real-World Application: Sound Waves

In audio engineering, two sound waves of different frequencies can be combined. The sum identity helps model the resultant wave pattern. For example, if a wave is described by $\sin(\omega_1 t)$ and another by $\sin(\omega_2 t)$, the product of these waves (which appears in amplitude modulation) can be expressed using sum-to-product identities derived from the sum and difference formulas. This allows engineers to design filters and

synthesize sounds with precision.

PRACTICE PROBLEMS: BUILD YOUR CONFIDENCE

The best way to internalize the **113 Sum And Difference Identities** is through consistent practice. Below are a few problems to try. Work through them step by step, referencing the formulas as needed.

1. Find the exact value of $\cos 105^\circ$ using sum or difference identities.
(Hint: $105^\circ = 60^\circ + 45^\circ$ or $135^\circ - 30^\circ$)
2. Simplify the expression:
 $\sin(40^\circ)\cos(20^\circ) + \cos(40^\circ)\sin(20^\circ)$.
3. Prove that $\tan(\pi/4 + x) = (1 + \tan x) / (1 - \tan x)$.

$$\begin{aligned} 4. \text{ Solve for } x \text{ in } [0, 2\pi): \sin(x + \pi/3) \\ = \sin(x - \pi/3). \end{aligned}$$

For the first problem, using $\cos(60^\circ + 45^\circ) = \cos 60^\circ \cos 45^\circ - \sin 60^\circ \sin 45^\circ = (1/2)(\sqrt{2}/2) - (\sqrt{3}/2)(\sqrt{2}/2) = (\sqrt{2}/4) - (\sqrt{6}/4) = (\sqrt{2} - \sqrt{6})/4$. This matches the known exact value.

CONCLUSION

The **113 Sum And Difference Identities** are a cornerstone of trigonometry. By understanding these six formulas—sine, cosine, and tangent for both sum and difference—you gain the ability to manipulate angles with confidence. They allow you to derive exact values, solve complex equations, and build bridges to more advanced concepts. While memorization helps, the

true power lies in recognizing the patterns and logic behind each identity. Practice regularly, use visualization on the unit circle, and connect the identities to real-world scenarios. As you master these tools, you will find that trigonometry becomes not only manageable but also deeply satisfying. Keep this guide handy as a reference, and soon you will be applying the sum and difference identities effortlessly in any mathematical context.